
Long-term research elucidates the spatial relationships

between imperiled bats & local fire management practices in the wake of White-nose Syndrome

Luke E. Dodd, Eastern Kentucky University, luke.dodd@eku.edu

### Introduction


- Insectivorous bats are important predators; studies suggest bats exert top-down pressure on insect prey communities & the ecosystem services of bats total billions of dollars.
- White-nose Syndrome (WNS) is an emergent disease responsible for the deaths of millions of bats across eastern North America. WNS has destabilized ecosystem community structure & relaxed niche partitioning.
- More than ever, effective management is needed for bat populations found across the fragmented forestlands of eastern North America.



# Objectives & Study Design

 We have elucidated the multi-trophic impacts of WNS in a fire-managed landscape.






# Methods & Base Results A - Access to excite deverter for state Line due for excitation forces A - Access to excite deverter for state Line of the excited force of the excited

## Derived Results & Implications

- Bats resistant to WNS (i.e., low-frequency echolocators) opt for less-cluttered conditions. As such, corridors are increasingly important as foraging habitats for bat communities at Mammoth Cave National Park (and more generally for eastern North America).
- Habitat 'quality' can be measured various ways; our work suggests the spatial dispersion of prey items
  across Mammoth Cave National Park does not necessarily correspond to the dispersion of nutrients
  across the landscape. Nutrition is a core concern for bats in the wake of WNS. Our work emphasizes
  the importance of assessing wildlife impacts beyond simple assessments of presence & abundance.



Biplot visualizing a distance-based redundancy analysis.

Red text represents acoustic abundance of bats. Arrows & black text represent constraining variables: fire presence & insect abundance. Ordination was significant (P < 0.05), with CAP1 and CAP2 accounting for 87% of explainable inertia in bat assemblage composition.

Map displaying the distribution of mean C:N ratios derived from Coleoptera, Diptera, Hymenoptera, & Lepidoptera captured during light-trapping surveys.

### Thanks!

- Funding from JFSP grants
   10-1-06-1 & 14-1-05-22.
- Technical support by
   C. Rose, J. Winters, &
   especially T. Culbertson!



# Coauthors & Affiliations

- M.B. Dickinson, US Forest Service, Northern Research Station
- M.J. Lacki, Univ. of Kentucky, Dept. of Forestry
- L.K. Rieske, Univ. of Kentucky, Dept. of Entomology
- N.S. Skowronski, US Forest Service, Northern Research Station
- S.C. Thomas, USNPS, Cumberland Piedmont Network
- R.S. Toomey, USNPS, Mammoth Cave National Park

